

Introduction to Scientific Computing Languages

Prof. **Paolo Bientinesi**

`pauldj@aices.rwth-aachen.de`

RWTHAACHEN
UNIVERSITY



Deutsche
Forschungsgemeinschaft

DFG

Fourier Transform

Let $f : \mathbf{R} \rightarrow \mathbf{C}$ be an integrable function.
Its *Fourier Transform* F is defined as the function

$$F(t) = \int_{-\infty}^{+\infty} f(x)e^{-2\pi ixt} dx.$$

Fourier transform: decomposition of a function into a series of waves.

Fourier Transform

Let $f : \mathbf{R} \rightarrow \mathbf{C}$ be an integrable function.
Its *Fourier Transform* F is defined as the function

$$F(t) = \int_{-\infty}^{+\infty} f(x)e^{-2\pi ixt} dx.$$

Fourier transform: decomposition of a function into a series of waves.

Discrete Fourier Transform

Let $\{x_k\}_0^{n-1}$ be a sequence of complex numbers.
Its *Discrete Fourier Transform* (DFT) is defined as the sequence

$$X_k = \sum_{j=0}^{n-1} x_j e^{-i2\pi \frac{k}{n} j}.$$

Fast Fourier Transform

One of the top-10 algorithms of the century

Definition

$$y := F_m x, \quad x, y \in \mathbb{R}^m$$

Fast Fourier Transform

One of the top-10 algorithms of the century

Definition

$$y := F_m x, \quad x, y \in \mathbb{R}^m$$

- $F_m = e^{\frac{-2\pi i}{m}kj}$, $k, j \in [0, 1, \dots, m-1]$
- $F_m = F_m^T$ (symmetry)
- Matlab: `fft`, `ifft`, `fftshift`, ...

Fast Fourier Transform

One of the top-10 algorithms of the century

Definition

$$y := F_m x, \quad x, y \in \mathbb{R}^m$$

- $F_m = e^{\frac{-2\pi i}{m}kj}$, $k, j \in [0, 1, \dots, m-1]$
- $F_m = F_m^T$ (symmetry)
- Matlab: `fft`, `ifft`, `fftshift`, ...

Recursive Definition

$$F_m = P_{m, \frac{m}{k}} (I_k \otimes F_{\frac{m}{k}}) D_{k, \frac{m}{k}} (F_k \otimes I_{\frac{m}{k}})$$

$$F_m = P_{m, \frac{m}{k}} (I_k \otimes F_{\frac{m}{k}}) D_{k, \frac{m}{k}} (F_k \otimes I_{\frac{m}{k}})$$

Identity matrix

$$I_k = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & & \\ \vdots & & \ddots & \\ 0 & & & 1 \end{bmatrix}_{[k \times k]}$$

$$F_m = P_{m, \frac{m}{k}} (I_k \otimes F_{\frac{m}{k}}) D_{k, \frac{m}{k}} (F_k \otimes I_{\frac{m}{k}})$$

Fourier matrix

$$F_m = e^{\frac{-2\pi i}{m}kj}, \quad k, j \in [0, 1, \dots, m-1]$$

Definitions

$$F_m = P_{m, \frac{m}{k}} (I_k \otimes F_{\frac{m}{k}}) D_{k, \frac{m}{k}} (F_k \otimes I_{\frac{m}{k}})$$

Kronecker Product

$$A = \begin{bmatrix} a_{00} & a_{01} & \dots \\ a_{10} & a_{11} & \\ \vdots & & \ddots \end{bmatrix}_{[m \times n]}, \quad A \otimes B = \begin{bmatrix} a_{00}B & a_{01}B & \dots \\ a_{10}B & a_{11}B & \\ \vdots & & \ddots \end{bmatrix}_{[? \times ?]}$$

$$F_m = P_{m, \frac{m}{k}} (I_k \otimes F_{\frac{m}{k}}) D_{k, \frac{m}{k}} (F_k \otimes I_{\frac{m}{k}})$$

Twiddle Factors

$$D_{k, \frac{m}{k}} = \begin{bmatrix} (D^0 = I) & 0 & \dots & 0 \\ 0 & D & & \\ & & D^2 & \\ \vdots & & & \ddots \\ 0 & & & & D^{k-1} \end{bmatrix}_{[m \times m]}$$

$$D = \text{diag}(e^{\frac{-2\pi i}{m} j}), \quad j \in [0, \dots, \frac{m}{k} - 1]$$

Definitions

$$F_m = P_{m, \frac{m}{k}} (I_k \otimes F_{\frac{m}{k}}) D_{k, \frac{m}{k}} (F_k \otimes I_{\frac{m}{k}})$$

Permutation

$$P_{m, \frac{m}{k}} = \text{stride}$$

$$P_{m, \frac{m}{2}} \left[\begin{array}{cccc|cccc} \times & \times & \times & \times & \dots & \times & \times & \\ \circ & \circ & \circ & \circ & \dots & \circ & \circ & \\ + & + & + & + & \dots & & & \\ \vdots & \vdots & & & & & & \\ \hline \hat{\times} & \hat{\times} & \hat{\times} & \dots & & & & \\ \hat{\circ} & \hat{\circ} & \hat{\circ} & \dots & & & & \\ \hat{+} & \hat{+} & \hat{+} & \dots & & & & \\ \vdots & \vdots & & & & & & \\ \vdots & \vdots & & & & & & \end{array} \right]_{[m \times m]} \rightarrow \left[\begin{array}{cccc|cccc} \times & \times & \times & \times & \dots & \times & \times & \\ \hat{\times} & \hat{\times} & \hat{\times} & \hat{\times} & \dots & \hat{\times} & \hat{\times} & \\ \hline \circ & \circ & \circ & \circ & \dots & \circ & \circ & \\ \hat{\circ} & \hat{\circ} & \hat{\circ} & \hat{\circ} & \dots & \hat{\circ} & \hat{\circ} & \\ \hline + & + & + & + & \dots & & & \\ \hat{+} & \hat{+} & \hat{+} & \hat{+} & \dots & & & \\ \hline \vdots & \vdots & & & & & & \\ \vdots & \vdots & & & & & & \end{array} \right]_{[m \times m]}$$

Definitions

$$F_m = P_{m, \frac{m}{k}} (I_k \otimes F_{\frac{m}{k}}) D_{k, \frac{m}{k}} (F_k \otimes I_{\frac{m}{k}})$$

Permutation

$$P_{12,3} \begin{bmatrix} \alpha_0 \\ \alpha_1 \\ \alpha_2 \\ \alpha_3 \\ \alpha_4 \\ \alpha_5 \\ \alpha_6 \\ \alpha_7 \\ \alpha_8 \\ \alpha_9 \\ \alpha_{10} \\ \alpha_{11} \end{bmatrix} \rightarrow \begin{bmatrix} \alpha_0 \\ \alpha_3 \\ \alpha_6 \\ \alpha_9 \\ \alpha_1 \\ \alpha_4 \\ \alpha_7 \\ \alpha_{10} \\ \alpha_2 \\ \alpha_5 \\ \alpha_8 \\ \alpha_{11} \end{bmatrix} \qquad P_{12,2} \begin{bmatrix} \alpha_0 \\ \alpha_1 \\ \alpha_2 \\ \alpha_3 \\ \alpha_4 \\ \alpha_5 \\ \alpha_6 \\ \alpha_7 \\ \alpha_8 \\ \alpha_9 \\ \alpha_{10} \\ \alpha_{11} \end{bmatrix} \rightarrow \begin{bmatrix} \alpha_0 \\ \alpha_2 \\ \alpha_4 \\ \alpha_6 \\ \alpha_8 \\ \alpha_{10} \\ \alpha_1 \\ \alpha_3 \\ \alpha_5 \\ \alpha_7 \\ \alpha_9 \\ \alpha_{11} \end{bmatrix}$$

Definition

$$Y := F_m X F_n$$

Definition

$$Y := F_m X F_n$$

- Sequence of 1d FFTs.

FFT of columns followed by FFT of rows, or viceversa

$$Y = (F_m X) F_n = F_m (X F_n)$$

Definition

$$Y := F_m X F_n$$

- Sequence of 1d FFTs.

FFT of columns followed by FFT of rows, or viceversa

$$Y = (F_m X) F_n = F_m (X F_n)$$

- 2-dimensional approach.

$$Y =$$

$$P_{m, \frac{m}{k}} (I_k \otimes F_{\frac{m}{k}}) D_{k, \frac{m}{k}} (F_k \otimes I_{\frac{m}{k}}) X (F_h \otimes I_{\frac{n}{h}}) D_{h, \frac{n}{h}} (I_h \otimes F_{\frac{n}{h}}) P_{n, \frac{n}{h}}^T$$

- Matlab: `fft2`, `ifft2`, ...